



11 Park Drive, Suite 12
Boston, MA 02215

Mouse soluble Tie1 recombinant protein

ORDERING INFORMATION

Catalog No: rAP-0272

Size: 2 µg; 10µg

Storage: <- 20° C

Synonyms:

Tyrosine kinase with immunoglobulin-like and EGF-like domains 1, JTK14, TIE, TIE1.

Introduction:

TIE-1 (tyrosine kinase with Ig and EGF homology domains 1) and TIE-2/Tek comprise a receptor tyrosine kinase (RTK) subfamily with unique structural characteristics: two immunoglobulin-like domains flanking three epidermal growth factor (EGF)-like domains and followed by three fibronectin type III-like repeats in the extracellular region and a split tyrosine kinase domain in the cytoplasmic region. These receptors are expressed primarily on endothelial and hematopoietic progenitor cells and play critical roles in angiogenesis, vasculogenesis and hematopoiesis. Human TIE-1 cDNA encodes a 1124 amino acid (aa) residue precursor protein with an 18 residue putative signal peptide, a 727 residue extracellular domain and a 354 residue cytoplasmic domain. Whereas two ligands have been described for TIE-2 [angiopoietin-1 (Ang1) and angiopoietin-2 (Ang2)], so far no ligand was found for TIE-1.

Description:

Soluble TIE-1 Mouse Recombinant fused with the Fc part of human IgG₁ produced in CHO is a monomeric, glycosylated, polypeptide containing 749 amino acids and having a total molecular mass of 260 kDa. Mouse TIE-1/Fc monomer has a calculated molecular mass of approximately 105 kDa. As a result of glycosylation, the recombinant protein migrates as an approximately 130 kDa protein in SDS-PAGE under reducing conditions.

The TIE1 Fc Chimera is purified by proprietary chromatographic techniques.

Source: CHO Cells

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

TIE-1 Fc Chimera was lyophilized from a concentrated (1 mg/ml) sterile solution containing 1x PBS.

Solubility:

It is recommended to reconstitute the lyophilized TIE-1 Fc Chimera in sterile water not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Stability:

Lyophilized sTIE-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TIE-1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Purity:

Greater than 90.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

Usage:

ProSpec's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com