



11 Park Drive, Suite 12  
Boston, MA 02215

## Human soluble VEGFR-2 FC recombinant protein

### ORDERING INFORMATION

Catalog No: rAP-0282

Size: 2 µg; 10µg

Storage: <- 20° C

**Synonyms:** KDR D1-7, sKDR D1-7, Kinase insert domain receptor, Protein-tyrosine kinase receptor Flk-1, CD309, type III receptor tyrosine kinase, FLK1, VEGFR-2.

#### Introduction:

Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes. All VEGF-receptors have seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. VEGFR-2 has a lower affinity for VEGF than the Flt-1 receptor, but a higher signaling activity. Mitogenic activity in endothelial cells is mainly mediated by VEGFR-2 leading to their proliferation. Differential splicing of the *flt-1* gene leads to the formation of a secreted, soluble variant of VEGFR-1 (sVEGFR-1). No naturally occurring, secreted forms of VEGFR-2 have so far been reported. The binding of VEGF<sub>165</sub> to VEGFR-2 is dependent on heparin.

**Description:** Soluble VEGFR-2 Fc Human Recombinant fused with the Fc part of human IgG<sub>1</sub> produced in baculovirus is a disulfide-linked homodimeric, glycosylated, polypeptide containing 757 amino acids and having a molecular mass of 160 kDa. The soluble receptor protein contains only the first 7 extracellular domains, which contain all the information necessary for ligand binding. The sKDR Fc Chimera is purified by proprietary chromatographic techniques.

**Source:** *Insect Cells*

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

#### Formulation:

KDR fusion protein was lyophilized from a concentrated (1 mg/ml) sterile solution containing 1x PBS pH-7.4.

**Solubility:** It is recommended to reconstitute the lyophilized VEGFR2 in sterile water not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

**Stability:** Lyophilized VEGFR-2 Fc/Chimera protein although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FLK1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). **Please prevent freeze-thaw cycles.**

**Purity:** Greater than 90.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

#### Biological Activity:

The activity of sVEGFR-2/Fc was determined by its ability to inhibit the VEGF-dependent proliferation of human umbilical vein endothelial cells. The ED<sub>50</sub> for this effect is typically 10-30 ng/ml.

**Usage:** Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)