



11 Park Drive, Suite 12
Boston, MA 02215

In vitro Human Lymphangiogenesis Tracing Kit (24-well plate format)

Ordering Information

Name of Product: In vitro Human Lymphangiogenesis Tracing Kit
Cat No: cAP-15

Introduction:

Lymphangiogenesis is a multistep process whereby new blood vessels develop from pre-existing lymphatic vasculature. Lymphoangiogenesis plays a key role in numerous physiological and pathological processes and understanding the mechanism of angiogenesis will therefore provide new approaches to the treatment of a wide range of pathologies.

Similar to vascular angiogenesis, lymphangiogenesis is also a complex process in which the following events are believed to play a critical role: (1) Proteolytic degradation of the extracellular matrix; (2) Directed migration of lymphatic endothelial cells; (3) Proliferation of lymphatic endothelial cells; (4) Deposition of new extracellular matrix; and (5) Formation of tubules and anastomosis of the newly formed lymphatic vessels.

The human LymphangioTracing Kit from Angio-Proteomie incorporates a proprietary system in which GFP-tagged human lymphatic endothelial cells are co-cultured with RFP-tagged human mesenchymal supporting cells in a specially designed medium. The lymphatic endothelial cells initially form small islands within the culture matrix. They subsequently begin to proliferate and then enter a migratory phase during which they move through the matrix to form threadlike tubule structures with lumens. These gradually join up (by 1 - 2 weeks) to form a network of anastomosing tubules which closely resembles the lymphatic capillary bed found in vivo.

The In vitro Human Lymphangiogenesis Tracing Kit (cAP-015) contains all of the materials necessary to perform multiple angiogenesis assays in a 24-well format. The Kit is designed that the testing materials, i.e. compounds, conditioned media, or tissue explants, can be added into the system at any time, ranging from the onset of lymphvasculogenesis to advanced lymphangiogenesis. The resulting effect on tubule formation (tubular length, number of branches et al) can be monitored throughout the whole process under inverted fluorescence microscope.

Reagents and Materials Provided

- (1) 1 x vial of mixture of GFP-tagged Human lymphatic ECs and RFP-tagged human mesenchymal supporting cells (-80°C or liquid N2)
- (2) 1 x 24-well Quick Coating Solution coated (cAP-01) plate (Room temperature)
- (3) 1 x 500ml of Endo-Growth Medium (cAP-02) (4°C)

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com

Protocols:

Day 1

1. Prewarm Endo-Growth Medium to 37°C in a water bath
2. Accurately pipette 24ml Endo-Growth Medium into a 50ml Falcon tube;
3. Rapidly thaw the vial of cryopreserved cells in a 37°C water bath;
4. Transfer all cells gently into 24ml pre warmed Endo-Growth medium;
5. Mix well the cells gently using a serological pipette;
6. Add 1.0ml of cell suspension to each well of the precoated 24-well plate.
7. Make sure the cells are evenly dispersed in the wells.
8. Place the plate in an incubator (37°C, 5% CO₂, and humidified).

Day 2


9. Take the plate from the incubator and examine cells under inverted fluorescence microscopy (GFP positive Human lymphatic ECs should sparsely and evenly distributed among RFP positive human mesenchymal supporting cells).
10. Wash the cells one with 2 ml of PBS
11. Add 2.0ml of fresh Endo-Growth medium (control) or Experimental media (Endo-Growth medium, plus pro- or anti-lymphangiogenic reagents according to customer's needs).
12. Place the plate back into the incubator.

Day 4, 6, 8, 10, 12, and 14.....

13. Replace the medium every 2 days until the end of the experiments.

(NOTE: Other commonly used endothelial growth medium may also been used with our human LymphangioTracing kit, but customer need optimize conditions before using those medium).

The images of lymphatic endothelial tubes can be monitored and recorded at any time during the whole experimental process under inverted fluorescence microscope.



Lymphatic capillary like tubes (Green) are shown in the above images after the cells were cultured with Endo-Growth Medium (cAP-02) for 12days. Tubules with lumens are indicated by arrows, human mesenchymal supporting cells (Red) are shown in the background.

This product is commercially produced for Research Use Only and under no circumstances that the product can be used for any human applications-20

Caution: Handling human derived products is potentially biohazardous. Although each cell strain has been tested negative for HIV, HBV and HCV DNA, diagnostic tests are not necessarily 100% accurate, therefore, proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working with these materials. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com