

## DATASHEET

### Fluorescent Antibody F(ab) Atto700

#### Anti-Rabbit IgG F(ab)Atto700

Goat anti-Rabbit IgG F(ab)Atto700

For Use in Research Only.

Not for Use in Human or Veterinary  
Diagnostic or Therapeutic Processes.

#### Kit Content (Cat. #: 2310-250UG)

250µg Anti-Rabbit IgG F(ab)Atto700 (H+L)  
1.0ml Glycerol buffer  
Product datasheet

#### Product Documentation

##### Anti-Rabbit IgG F(ab)Atto700

Anti-Rabbit IgG F(ab)Atto700 is the antigen-specific fragment of the antibody obtained by papain cleavage. Affinity purification removed essentially all goat serum proteins, including immunoglobulins not specifically binding to rabbit IgG. F(ab) fragment was purified by ion exchange chromatography (IEX). Anti-Rabbit IgG F(ab) is conjugated to Atto700 NHS (Abs.max. 700 nm; Em.max. 719 nm) with a degree of labelling of 1-3 and subsequent purification by gel filtration. The antibody fragment is supplied in unit sizes of 250µg.

#### Reconstitution of Antibodies with Glycerol-PBS

Add 125µl Glycerol buffer to the lyophilized secondary antibody to reconstitute a 2mg/ml stock solution. Vortex for 10sec until completely dissolved. Final concentrations of the antibody buffer: 0.01M sodium phosphate, 0.1M NaCl, pH 7.4, 5mM NaN<sub>3</sub> in 50% glycerol.

#### Working Dilution

Each individual user should determine the optimum working dilution empirically for the systems. Dilutions of 1:300 – 1:500 are suited for many applications.

#### Determining the Degree of Labeling (DOL)

##### 1. Protein Concentration

Determination of the protein concentration by UV absorption measurement at 280nm ( $\epsilon_{\text{max}} = 203,000 \text{ M}^{-1}\text{cm}^{-1}$ ).

1 OF 2

*For product inquiries please contact:*

cusserv@hypermol.com  
Fon: +49 (0)521 9876228  
Fax: +49 (0)521 9876231  
www.hypermol.com

## DATASHEET

---

### Fluorescent Antibody F(ab) Atto700

#### 2. Degree of Labelling

The degree of labeling (DOL or dye/protein ratio) is usually determined by absorption spectroscopy making use of the Lambert-Beer law: Absorbance (A) = extinction coefficient ( $\epsilon$ ) × molar concentration × path length (d). Simply measure the UV-VIS spectrum of the conjugate in solution in a quartz cuvette. Dilute the solution, if necessary to measure within the linear range.

$$\text{DOL} = \frac{A_{700} \cdot 203,000}{A_{280} - (A_{700} \cdot 0.41) \cdot 120,000}$$

$A_{700}$  = maximal absorbance at 700nm measured in a cuvette with a pathlength of 1 cm.

$A_{280}$  = maximal absorbance at 280nm measured in a cuvette with a pathlength of 1 cm.

203,000 = molar extinction coefficient ( $\epsilon$ ) at the longest-wavelength absorption maximum ( $\text{M}^{-1}\text{cm}^{-1}$ ).

120,000 = molar extinction coefficient ( $\epsilon$ ) at the longest-wavelength absorption maximum ( $\text{M}^{-1}\text{cm}^{-1}$ ).

0.41 = correction factor for the fluorophore's absorbance at 280nm.

#### Storage and Stability

For continuous use, store at 2-8 °C for up to three months. For extended storage, the solution may be frozen in working aliquots at -20 °C. Frozen aliquots are stable for at least six months. Avoid repeated freeze/thawing. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Protect fluorescent conjugates from light.

HYPERMOL EK WARRANTS THAT ITS PRODUCT ARE CONFORM TO THE INFORMATION CONTAINED IN THIS PUBLICATION. THE PURCHASERS MUST DETERMINE THE SUITABILITY OF THE PRODUCT(S) FOR THEIR PARTICULAR USE. ADDITIONAL TERMS AND CONDITIONS MAY APPLY (SEE REVERSE SIDE OF THE INVOICE). HYPERMOL BRANDS ARE SOLD EXCLUSIVELY BY HYPERMOL EK, GERMANY AND AUTHORIZED DISTRIBUTORS.

*For product inquiries please contact:*

cusserv@hypermol.com  
Fon: +49 (0)521 9876228  
Fax: +49 (0)521 9876231  
www.hypermol.com