

DATASHEET

Actin

From porcine skeletal muscle
(α -skeletal muscle actin, >99% purity)

For Use in Research Only.
Not for Use in Diagnostic Processes.

Quantity: 1x1.0mg / 10x1.0mg
Cat. #: 8111-01

Product Description

G-actin from porcine skeletal muscle is a single chain polypeptide with a molecular weight of 42kD consisting of 375 amino acids. G-actin (1mg) is supplied as lyophilized powder containing 2mM Tris-Cl pH 8.2, 0.4mM ATP, 0.1mM DTT, 0.08mM CaCl₂ and 0.2% disaccharides, when reconstituted with 1ml H₂O. For dilution of G-actin or exchange of ligand buffer into actin compatible buffer, MonoMix (Cat no.: 5100-0*) may be used. In the absence of nucleators, the polymerization of G-actin is initiated by PolyMix (Cat no.: 5000-0*).

Product Handling

Preparation of a stock solution

Add e.g. 1ml of H₂O to the tube with G-actin to obtain a stock solution of 1mg/ml (23.8 μ M). A stock concentration of 4mg/ml (95.2 μ M) should not be exceeded to avoid spontaneous polymerization. Allow the G-actin solution to rehydrate for 2min at room temperature and dissolve using a pipette. If G-actin is used for polymerization with PolyMix (Cat no.: 5000-0*), we recommend to prepare a G-actin stock of 1.1mg/ml (26.1 μ M), as addition of PolyMix will dilute the G-actin/F-actin stock to a final concentration of 1mg/ml (23.8 μ M).

The actin stock is ready for use. For critical assays we recommend gel filtration or dialysis against 50-100vol. of MonoMix O/N and subsequent centrifugation at 100.000xg for 60min to remove any aggregates from freeze-drying. For less critical assays the actin working stock may be pre-spun (15.000xg, 10min).

Product Storage and Stability

Freeze-dried G-actin stored at -70°C upon arrival will be stable in performance for at least 6 months from the date of purchase. Solubilized G-actin is kept on ice and should be used within 1 week. Polymerized actin (F-actin) kept on ice is stable for at least 2 weeks. Avoid refreezing of G-actin. Never refreeze F-actin solutions.

HYPERMOL EK WARRANTS THAT ITS PRODUCT ARE CONFORM TO THE INFORMATION CONTAINED IN THIS PUBLICATION. THE PURCHASERS MUST DETERMINE THE SUITABILITY OF THE PRODUCT(S) FOR THEIR PARTICULAR USE. ADDITIONAL TERMS AND CONDITIONS MAY APPLY (SEE REVERSE SIDE OF THE INVOICE). HYPERMOL BRANDS ARE SOLD EXCLUSIVELY BY HYPERMOL EK, GERMANY AND AUTHORIZED DISTRIBUTORS.

For product inquiries please contact:

cusserv@hypermol.com
Fon: +49 (0)521 9876228
Fax: +49 (0)521 9876231
www.hypermol.com