
DATA SHEET

CyAL-5 2-Deoxglucose Optical probe

For Imaging multiple tumor types such as breast, glioblastoma, colon and prostate

Product Description: CyAL-5 2-deoxyglucose (CyAI-5 2-DG) is a fluorescent imaging agent designed to target a wide spectrum of cancers such as breast, glioblastoma, colon and prostate in vitro as well as in vivo. 2-DG is a glucose analog that utilizes the GLUT transporters for entry into the cell. One of the hallmarks of many cancer cells is an elevated uptake of glucose. Numerous fluorophores, including 800CW [1] and 2-NBDG [2] have been used to label 2-DG for tumor detection. FDG has been used extensively in PET imaging for diagnosis and tumor response monitoring in various types of cancer [3,4].



The recommended individual dose per mouse will range from 10-20 nmol, depending upon tumor type, size and location. Material should be reconstituted in water or PBS.

References:

[1] Kovar JL, Volcheck W, Olive DM, Simpson MA (2007). Abstract #5527, Poster presentation, AACR Annual Meeting [2] Lloyd PG, Hardin CD, Sturek M(1999). Physiol Res 48:401-410.

[3] Hicks RJ, Wahl RL (2010).PET diagnosis and response monitoring in oncology, Molecular Imaging, Principles and Practice, 875-895.

[4] Mankoff DA(2010). PET imaging in cancer clinical trials, Molecular Imaging, Principles and Practice, p1179-1191.

For further information or to place an order please contact Dr. Brian Gray by email: briangray@mtarget.com or phone: 610-738-7938

Storage and Handling: Upon receipt, store at -20°C prior to reconstitution. Reconstituted material should be used within two weeks.

CyAL-5 is sold under an exclusive license from Harvard Medical School and Massachusetts General Hospital.

MOLECULAR TARGETING TECHNOLOGIES, INC.



833 Lincoln Ave., Unit 9 West Chester, PA 19380 P: 610.738.7938 F: 610.738.7928 Contact us: info@mtarget.com; website: www.mtarget.com