



11 Park Drive, Suite 12  
Boston, MA 02215

## Human Insulin-Like Growth Factor 1 (IGF1)

### ORDERING INFORMATION

Catalog No: rAP-0043;  
Size: 20 µg; 100 µg  
Storage: <- 20° C

### Synonyms:

Somatomedin C, IGF-I, IGF1, IGF1, IGF-IA, Mechano growth factor, MGF.

### Introduction:

The somatomedins, or insulin-like growth factors (IGFs), comprise a family of peptides that play important roles in mammalian growth and development. IGF1 mediates many of the growth-promoting effects of growth hormone (GH; MIM 139250). Early studies showed that growth hormone did not directly stimulate the incorporation of sulfate into cartilage, but rather acted through a serum factor, termed 'sulfation factor,' which later became known as 'somatomedin' (Daughaday et al., 1972). Three main somatomedins have been characterized: somatomedin C (IGF1), somatomedin A (IGF2; MIM 147470), and somatomedin B (MIM 193190) (Rotwein, 1986; Rosenfeld, 2003).

### Description:

Insulin-Like Growth Factor-I Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 70 amino acids and having a molecular mass of 7655 Dalton. IGF-I is purified by proprietary chromatographic techniques.

### Source:

*Escherichia Coli.*

### Physical Appearance:

Sterile Filtered White Lyophilized (freeze-dried) powder.

### Formulation:

The protein was lyophilized after extensive dialysis against 50mM acetic acid.

### Solubility:

It is recommended to reconstitute the lyophilized IGF-1 in sterile 18MΩ-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

### Stability:

Lyophilized Insulin-Like Growth Factor-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IGF1 should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

### Purity:

Greater than 98.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

### Amino acid sequence:

The sequence of the first five N-terminal amino acids was determined and was found to be Gly-Pro-Glu-Thr-Leu. N-terminal methionine has been completely removed enzymatically.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)



11 Park Drive, Suite 12  
Boston, MA 02215

**Biological Activity:**

The ED<sub>50</sub>, calculated by the dose-dependant proliferation of murine BALB\C 3T3 cells (measured by <sup>3</sup>H-thymidine uptake) is < 1.0 ng/ml, corresponding to a specific activity of 1x10<sup>6</sup> U/mg.  
For most *in-vitro* applications, IGF-I exerts its biological activity in the concentration range of 0.2-20 ng/ml.

**Protein content:**

Protein quantitation was carried out by two independent methods:

1. UV spectroscopy at 280 nm using the absorbency value of 0.502 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics).
2. Analysis by RP-HPLC, using a calibrated solution of IGF1 as a Reference Standard

**Usage:**

Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.