



11 Park Drive, Suite 12  
Boston, MA 02215

## Human Monocyte Chemotactic Protein-4 (MCP-4)

### ORDERING INFORMATION

Catalog No: rAP-0103;

Size: 2 µg; 10 µg

Storage: <- 20° C

### Synonyms:

Small inducible cytokine A13, CCL13, Monocyte chemotactic protein 4, MCP-4, Monocyte chemoattractant protein 4, CK-beta-10, NCC-1, chemokine (C-C motif) ligand 13, NCC1, CKb10, SCYL1, SCYA13, MGC17134.

### Introduction:

Chemokine (C-C motif) ligand 13 (CCL13) is a small cytokine belonging to the CC chemokine family. Its gene is located on human chromosome 17 within a large cluster of other CC chemokines. CCL13 induces chemotaxis in monocytes, eosinophils, T lymphocytes, and basophils by binding cell surface G-protein linked chemokine receptors such as CCR2, CCR3 and CCR5. Activity of this chemokine has been implicated in allergic reactions such as asthma. CCL13 can be induced by the inflammatory cytokines interleukin-1 and TNF-α.

### Description:

Monocyte Chemotactic Protein-4 Human Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 74 amino acids and having a molecular mass of 8510 Dalton. The MCP-4 is purified by proprietary chromatographic techniques.

### Source:

*Escherichia Coli.*

### Physical Appearance:

Sterile Filtered White Lyophilized (freeze-dried) powder.

### Formulation:

The CCL13 protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives.

### Solubility:

It is recommended to reconstitute the lyophilized Monocyte Chemotactic Protein-4 in sterile 18MΩ-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

### Stability:

Lyophilized MCP4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL13 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Please prevent freeze-thaw cycles.**

### Purity:

Greater than 95.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

### Amino acid sequence:

The sequence of the first five N-terminal amino acids was determined and was found to be Gln-Pro-Asp-Ala-Leu.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)



11 Park Drive, Suite 12  
Boston, MA 02215

**Biological Activity:**

The specific activity as determined by the ability of MCP-4 to chemoattract human monocytes at 10-100 ng/ml.

**Usage:**

Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, [angioproteomie@gmail.com](mailto:angioproteomie@gmail.com)