



11 Park Drive, Suite 12
Boston, MA 02215

Human Tumor Necrosis Factor-Alpha (TNF- α), His Tag

ORDERING INFORMATION

Catalog No: rAP-0159;

Size: 10 μ g; 50 μ g

Storage: <- 20° C

Synonyms:

TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNF-a, Cachectin, DIF, TNFA, TNFSF2.

Introduction:

Tumor necrosis factor is a cytokine involved in systemic inflammation and is a member of a group of cytokines that all stimulate the acute phase reaction. TNF is mainly secreted by macrophages. TNF causes apoptotic cell death, cellular proliferation, differentiation, inflammation, tumorigenesis and viral replication, TNF is also involved in lipid metabolism, and coagulation. TNF's primary role is in the regulation of immune cells.

Dysregulation and, in particular, overproduction of TNF have been implicated in a variety of human diseases- autoimmune diseases, insulin resistance, and cancer.

Description:

Tumor Necrosis Factor- α Human Recombinant His produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 157 amino acids fragment (77-233) and having a molecular mass of 21.85 kDa with an amino-terminal hexahistidine tag.

The TNF-alpha His is purified by standard chromatographic techniques.

Source:

Escherichia Coli.

Physical Appearance:

Sterile Filtered clear solution.

Formulation:

TNF-a His is supplied in 10mM Tris, 100mM NaCl, 50% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time.

Please avoid freeze thaw cycles.

Purity:

Greater than 95.0% as determined by:

(a) Analysis by RP-HPLC.

(b) Analysis by SDS-PAGE.

Usage:

Angio-Proteomie's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com