

DATASHEET

Rhodamine-Actin

Rhodamine-Actin

From rabbit skeletal muscle
(α -skeletal muscle actin)

For Use in Research Only.
Not for Use in Diagnostic Processes.

Quantity: 1x100 μ g
Cat.#: 8103-01

Product Description

G-actin from rabbit skeletal muscle is a single chain polypeptide with a molecular weight of 42kD consisting of 374 amino acids. Rhodamine-Actin is chemically modified G-actin by coupling of tetramethylrhodamine-5/6-isothiocyanate to Cys374 at the C-term of the actin molecule. The dye/protein molar ratio of Rhodamine-Actin is >0.5:1, using a molar extinction coefficient of 72,000 cm⁻¹ M⁻¹ at 558nm. The protein concentration of Rhodamine-Actin was determined by Bradford method. Rhodamine-Actin is supplied as a lyophilized powder containing 5mM Tris-Cl pH 8.2, 1mM ATP, 1mM DTT and 0.2mM CaCl₂, when reconstituted with H₂O to a 1mg/ml solution.

For dilution of G-actin or exchange of ligand buffer into actin compatible buffer, MonoMix (Cat no.: 5100-0*) may be used. In the absence of nucleators, the polymerization of G-actin can be initiated by PolyMix (Cat no.: 5000-0*).

Product Handling

Preparation of a working stock

Example: Add 100 μ l of H₂O to a tube with Rhodamine-Actin to obtain a working stock of 1mg/ml (24 μ M) and vortex for 30secs. The final concentration of G-actin should not exceed 3mg/ml. Allow the G-actin solution to rehydrate for 5min at room temperature, dissolve the pellet by pipetting several times up and down to obtain a homogeneous, pinkish Rhodamine-Actin solution. Leave to fully rehydrate for another 2 min, vortex for 30secs and spin 1min, 15.000xg. For standard applications the Rhodamine-Actin working stock is now ready to use. For critical assays the working stock should at least be dialyzed (100vol. of MonoMix, O/N) followed by high-speed centrifugation. For less critical assays the actin working stock may be filtered (0.2-0.45 μ m) or pre-spun (15.000rpm, 10min).

Product Storage and Stability

The highest product performance is achieved with Rhodamine-Actin stored at -70° until used. Once dissolved Rhodamine-Actin is kept on ice protected from light and will be stable for 5 days. Avoid re-freezing of solubilized Rhodamine-Actin.

HYPERMOL EK WARRANTS THAT ITS PRODUCT ARE CONFORM TO THE INFORMATION CONTAINED IN THIS PUBLICATION. THE PURCHASERS MUST DETERMINE THE SUITABILITY OF THE PRODUCT(S) FOR THEIR PARTICULAR USE. ADDITIONAL TERMS AND CONDITIONS MAY APPLY (SEE REVERSE SIDE OF THE INVOICE). HYPERMOL BRANDS ARE SOLD EXCLUSIVELY BY HYPERMOL EK, GERMANY AND AUTHORIZED DISTRIBUTORS.

1 of 1

For product inquiries please contact:

cusserv@hypermol.com
Fon: +49 (0)521 9876228
Fax: +49 (0)521 9876231
www.hypermol.com